

Retrieval of Lodged Foreign Body from the Root Canal - A Case Report

¹Dr. Damini Patil, Post Graduate Student, Department of Conservative Dentistry and Endodontics, MIDSR Dental college, Latur.

²Dr. Ramchandra Kabir, Reader, Department of Conservative Dentistry and Endodontics, MIDSR Dental college, Latur.

³Dr. Sunanda Gaddalay, Professor & HOD, Department of Conservative Dentistry and Endodontics, MIDSR Dental college, Latur.

Corresponding Author: Dr. Damini Patil, Post Graduate Student, Department of Conservative Dentistry and Endodontics, MIDSR Dental college, Latur.

Citation of this Article: Dr. Damini Patil, Dr. Ramchandra Kabir, Dr. Sunanda Gaddalay, “Retrieval of Lodged Foreign Body from the Root Canal - A Case Report”, IJDSIR- June - 2022, Vol. – 5, Issue - 3, P. No. 254 – 258.

Copyright: © 2022, Dr. Damini Patil, et al. This is an open access journal and article distributed under the terms of the creative commons attribution non-commercial License. Which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Type of Publication: Case Report

Conflicts of Interest: Nil

Abstract

Accidental ingestion or aspiration of a variety of foreign bodies is a common endodontic emergency. A foreign body in the tooth, however, is rare. Retrieval of foreign objects from the teeth is a challenging aspect in clinical practice.

A detailed case history, including history of habitual placement of the objects, clinical and radiographic examinations are necessary to come to a conclusion about the nature, size, location of the foreign body and the difficulty involved in its retrieval. Sometimes the foreign objects fracture inside the tooth while exploration by the patient. These foreign objects may act as a potential source of infection and may later lead to complications. This paper discusses a case of foreign object impaction in the teeth, the possible aetiology along with its management.

Keywords: Foreign objects, Habitual placement, non-surgical technique, Source of infection, Trauma

Introduction

Presence of foreign objects in root canal is one of the troublesome incidents in endodontic therapy. The chance of these foreign objects getting impacted into the tooth is more when pulp chamber is open either because of traumatic injury or large carious exposure.

These foreign objects can be easily retrieved if they are located within the pulp chamber, but once the object has been pushed apically, their retrieval may be complicated. Apical surgical procedures may sometimes be unavoidable.¹

The following case describes a foreign object impacted into the apical third of a maxillary central incisor, which was retrieved by simple nonsurgical intracanal means.

Case Report 1

A 28-year-old Male reported to the Department of Conservative Dentistry and Endodontics with a complaint of pain and also removal of stapler pin in the root canal of maxillary central incisor. Patient admitted of placing stapler pin inside the root canal to remove food plugs from the teeth. However, the stapler pin got lodged accidentally in the root canal of central incisor. Patient tried to remove the stapler pin with a needle and was unsuccessful.

On examination, the pulp chamber was found to be open to oral cavity but was occluded with food plugs. Radiographic examination revealed the presence of a radio-opaque object in the root canal extending from middle third to the apex of the root [Figure 1]. It was decided to retrieve the stapler pin by nonsurgical technique, and thereafter, complete the routine endodontic treatment in all the two teeth.

Conventional access cavity was refined to facilitate access for instrumentation. An ISO no. 20 K-file (DENTSPLY Maillefer, Ballaigues, Switzerland) was used to bypass the stapler pin. Retrieval was done by attempting to engage the stapler pin between ISO no. 20 H-file (DENTSPLY Maillefer) and canal wall then pulling it out coronally, which was then grasped with tweezers and was retrieved. The retrieved stapler pin was six millimeters in length [Figure 2].

The working length was confirmed using an apex locator (Root ZX; J Morita, Tokyo, Japan) as well as radiographically using a stainless-steel K-file no. 15 [Figure-3]. The root canals were cleaned and shaped using Neo hybrid rotary file system (Neo Endo, Orikam, Health Care Solution, India) in a crown down motion. Three percent sodium hypochlorite (Prime Dental, India), 17% ethylenediamine tetra acetic acid liquid (EDTA) (Dent wash, Prime Dental, India), and isotonic

saline was used as irrigants. An intracanal calcium hydroxide medicament was placed [Figure-4].

Patient was recalled after 3 weeks for the follow-up. At subsequent appointment, canal was irrigated with 2.5% NaOCl (Prime Dental, India) and 2% chlorhexidine (Chlor X, Prevest Denpro). The canal was dried with paper points and Bio dentin was placed with pluggers until thickness of 6 mm [Figure-4].

A wet cotton pellet was placed in the canal and access cavity was sealed with temporary cement. In next appointment, root canal was obturated with GP using lateral condensation technique using AH plus sealer (Dentsply, Maillefer, Konstanz, Germany) [Figure 5]. Access cavity sealed with glass ionomer cement [Figure-6]. On a follow-up examination after three months, the teeth were asymptomatic.

Discussion

A number of foreign objects were reported to be lodged in the pulp chamber or root canals of the tooth, which ranged from stapler pin², pencil leads³, darning needles⁴, metal screws⁵, beads⁶, plastic chop stick⁷, hat pins⁸, dress maker pins⁹, conical metallic object¹¹.

During emergency root canal treatment, the patient remains in the office with a draining tooth for an hour or even more and finally ending the appointment by sealing the access cavity¹². With the access cavity closed, new strains of microorganism systems and foreign bodies can be prevented from entering the root canal¹³.

Patient should be well educated regarding the phases of root canal treatment and the importance of completion of treatment, to avoid undue consequences.

A radiograph can be of diagnostic significance especially if the foreign body is radio-opaque. Specialized radiographic techniques¹.

such as Radio visiography, three-dimensional CAT (Computerized axial tomography) scans can play a

pivotal role in the localization of these foreign objects inside the root canal.

Foreign objects in root canals can act as focus of infection. Complications can follow if these impacted foci of infection are not eliminated at the right time. Actinomycosis following placement of piece of jewellery chain into a maxillary central incisor¹⁴, and chronic maxillary sinusitis of dental origin developed due to pushing of foreign bodies into the maxillary sinus¹⁵, has been reported. Hence, prompt attempts at their retrieval should be initiated.

For retrieval of foreign objects lying in the pulp chamber or canal using ultrasonic instruments¹⁶, the Masserann kit¹⁷, modified Castroviejo needle holders¹⁸, the Steglitz forceps¹⁹, have been used. EDTA has been suggested as a useful aid in lubricating the canal when attempting to remove the foreign object¹⁹.

The use of an operating microscope along with ultrasonic²⁰ provides the clinician to visualize any intraradicular metallic obstructions.

Nonetheless, retrieval of the object may be difficult when it is lodged in periapical region. Periapical surgery or intentional reimplantation²¹ should be considered to remove such objects.

In this case stapler pin was located within the root canal and confirmed by diagnostic radiograph. Access to the foreign body was improved by flaring the canal²² coronally. Every attempt has to be made to bypass and retrieve the stapler pin by pulling it out coronally.

If foreign objects are found in the root canal, prompt, but cautious attempts should be made to retrieve it first by simple nonsurgical means. Finally, when the foreign object resists all efforts for removal a surgical procedure may be the only viable alternative.

Figures



Figure no 1: Pre-Operative Radiograph



Figure no 2: Retrieval of stapler pin from the canal radiograph



Figure no 3: Working length Radiograph



Figure 5: Bio dentin apical plug placed using hand pluggers



Figure 4: Calcium Hydroxide as an intracanal medicament placed within the canal



Figure no 2 A: Post Obturation Radiograph

References

1. Prabhakar AR, Namineni S, Subhadra HN. Foreign body in the apical portion of a root canal in a tooth with an immature apex: A case report. *Int Endod J* 2008; 41:920-7.
2. McAuliffe N, Drager NA, Hunter B. Staple diet: A foreign body in a tooth. *Int J Paediatr Dent* 2005; 15:468-71.
3. Hall JB. Endodontics – Patient performed. *J Dent Child* 1969; 36:213-5.
4. Nernst H. Foreign body in a root canal. *Quintessence Int*
5. Prabhakar AR, Basappa N, Raju OS. Foreign body in a mandibular permanent molar - a case report. *J Indian Soc Pedod Prev Dent*, 1998;16:120-1.
6. Reddy VV, Mehtha DS. Beads. *Oral Surg Oral Med Oral Pathol*, 1990;69:769-70.
7. Toida M, Ichihara H, Oku Tomi T, Nakamura K, Ishimaru JI. An unusual foreign body in an unerupted supernumerary tooth. *Br Dent J*, 1992; 173:345-6.
8. Zillich RM, Pickens TN. Patient induced blockage of the root canal. Report of a case. *Oral Surg Oral Med Oral Pathol* 1982; 54:689-90.
9. Turner CH. An unusual foreign body. *Oral Surg Oral Med Oral Pathol* 1983; 56:226.
10. Gelfman WE, Cheris LJ, Williams AC. Self-attempted endodontics- a case report. *J Dent Child* 1969; 36:283-4.
11. Lamster IB, Barenie JT. Foreign objects in the root canal. *Oral Surg Oral Med Oral Pathol* 1977; 44:483-6.
12. Weine FS. Endodontic Emergency treatment. In: Weine FS, editor. *Endodontic therapy*. St Louis: Mosby; 2004. p. 72-103.
13. Nair PN. On the causes of persistent apical periodontitis: A review. *Int Endod J* 2006; 39:249-81.
14. Goldstein BH, Scuibba JJ, Laskin DM. Actinomycosis of the maxilla: A review of literature and report of a case. *J Oral Surg* 1972; 3:362-6.
15. Costa F, Robiony M, Toro C, Sembronio S, Politi M. Endoscopically assisted procedure for the removal of a foreign body from the maxillary sinus and contemporary endodontic surgical treatment of the tooth. *Head Face Med* 2006; 2:37.
16. Meindinger DL, Kabes BJ. Foreign object removal utilizing the cavi- endo ultrasonic instrument. *J Endod* 1985; 11:301-4.
17. Williams VD, Bjorndal AM. The Masserann technique for the removal of fractured posts in endodontically treated teeth. *J Prosthet Dent* 1983; 49:46-8.
18. Fros UG, Berg JO. A method for the removal of broken endodontic instruments from root canals. *J Endod* 1983; 9:156-9.
19. Lumley PJ, Walmsley AD. The removal of foreign objects from root canals. *Dent update* 1990; 17:420-3.
20. Nehme WB. Elimination of intracanal metallic obstructions by abrasion using operational microscope and ultrasonics. *J Endod* 2001; 27:365-7.
21. Srivastava N, Vineeta N. Foreign body in the periradicular area. *J Endod* 2001; 27:593-4.
22. McCulloch AJ. The removal of restorations and foreign objects from root canals. *Quintessence Int* 1993; 24:245-9.